

Images & turbulence — 08

The wavefront is, modulo $\lambda/2\pi$, proportional to the phase $\Phi(r)$ of the wave $\Psi(r)$ which has went through the turbulent atmosphere before reaching the telescope:


$$\Psi(\vec{r}) = A(\vec{r}) \exp\{i\Phi(\vec{r})\}$$

Note that this phase can be decomposed following a base of polynomials, for example Zernike ones:


$$\Phi(\vec{r}) = \sum_i a_i Z_i(\vec{r})$$

Images & turbulence — 09

polynômes de Zernike



entrance pupil



Images & turbulence – 10

turbulence intensity [$m^{1/3}$]

$$r_0 = 0.185 \lambda^{6/5} \cos(\gamma)^{3/5} \left[\int_0^\infty C_n^2(z) dz \right]^{-3/5}$$

dimension of r_0 ? value in band H knowing r_0 at 500nm (10cm) ?...

$$\tau_0 = 0.36 \frac{r_0}{\bar{v}}$$

$$\epsilon_0 = 0.98 \frac{\lambda}{r_0}$$

$$\theta_0 = 0.314 \frac{r_0}{\bar{h}}$$

$$\bar{v} = \left(\frac{\int C_n^2(h) v(h)^{5/3} dh}{\int C_n^2(h) dh} \right)^{3/5}$$

$$N_s \simeq 0.34 \left(\frac{D}{r_0} \right)^2$$

$$\bar{h} = \left(\frac{\int C_n^2(h) h^{5/3} dh}{\int C_n^2(h) dh} \right)^{3/5}$$

Number of speckles for $r_0=10\text{cm}$ and $D=1\text{m}$?...

Images & turbulence – 11

r_0 in band H knowing r_0 at 500nm ?...

$$r_0 = 0.185 \lambda^{6/5} \cos(\gamma)^{3/5} \left[\int_0^\infty C_n^2(z) dz \right]^{-5/3}$$

$$r_0^{H=1.65 \mu\text{m}} = r_0^{500 \text{ nm}} \left(\frac{1.65}{0.5} \right)^{6/5} \simeq 0.42$$

Number of speckles for $r_0=10\text{cm}$ and $D=1\text{m}$?...

$$N_S^{500 \text{ nm}} \simeq 0.34 \left(\frac{1.0}{0.1} \right)^2 \simeq 34$$

$$N_S^H \simeq 0.34 \left(\frac{1.0}{0.42} \right)^2 \simeq 2$$

Images & turbulence – 12

$$\Phi_{\varphi}(\vec{\nu}) = 0.0228 r_0^{-\frac{5}{3}} \left(\nu^2 + \frac{1}{\mathcal{L}_0^2} \right)^{-\frac{11}{6}}$$


Power Spectral Density (PSD) of the phase, function of the spatial frequency

Kolmogorov/von Kármán model

- Kolmogorov : outerscale of turbulence \mathcal{L}_0 is infinite.
- One can refine the model by considering also ℓ_0 .
- \exists other models with a finite \mathcal{L}_0 (and a non-zero ℓ_0).

Images & turbulence – 13

$$\Phi_{\varphi}(\vec{\nu}) = 0.0228 r_0^{-\frac{5}{3}} \left(\nu^2 + \frac{1}{\mathcal{L}_0^2} \right)^{-\frac{11}{6}}$$




Energy cascade:

wind shear \Rightarrow turbulent energy injected into the system via a large eddy (\mathcal{L}_0) which splits into smaller and smaller eddies (l_0), and is finally viscously dissipated.

Interval $[l_0, \mathcal{L}_0]$ = inertial range.

(A reminder of discrete Fourier transform (DFT)...)



Images & turbulence – 14

$$\Phi_{\varphi}(\vec{\nu}) = 0.0228 r_0^{-\frac{5}{3}} \left(\nu^2 + \frac{1}{L_0^2} \right)^{-\frac{11}{6}}$$

Which, numerically written, and by considering wavefronts made of 'dim' pixels corresponding to 'L' meters, becomes:
(re-writing - "de-dimensionalizing" - the equation with $L_0=L_0 L/L$ and $\nu=\nu L/L\dots$)

```
freq = findgen(dim)
dsp   = .0228*(L/r0)^(5/3.)*L^2*(freq^2+(L/L0)^2)^(-11./6)
```

And which (with the right frequency scale) can be plot with:

```
plot_oo, 1./L*findgen(dim), dsp, XR=[1/L/1.2,dim*1/L*1.2], /XS
```

=> make a function that computes $\text{PSD}(L_0, r_0, \text{dim}, L)$ and plot it for different $[r_0, L_0] \dots$ [with, for example: $\text{dim}=1000, L=100., r_0=0.1, L_0=100.,10.,1.]$

(IDL – 2)

Example of a function that computes the sum of two parameters:

```
function sum_of_two_parameters, par1, par2  
result=par1+par2  
return, result  
end
```

Compile and run the function (written, e.g., in a file sum2.pro):

```
idl > .r sum2  
→ % Compiled module: SUM_OF_TWO_PARAMETERS  
idl > res = sum_of_two_parameters(2,1)  
idl > print, res  
→ 3
```