

# Imaging through turbulence

(image formation, atmospheric turbulence, intro to adaptive optics)

Marcel Carbillet  
[[marcel.carbillet@oca.eu](mailto:marcel.carbillet@oca.eu)]


[lagrange.oca.eu/carbillet/enseignement/M1-MAUCA/](http://lagrange.oca.eu/carbillet/enseignement/M1-MAUCA/)

# Content

- High-angular resolution imaging in astronomy
- Atmospheric turbulence (reminder)
- Numerical modelling of perturbed wavefronts
- Formation of resulting images (+detection noises)
- *(Introduction to speckle interferometry)*
- Introduction to adaptive optics (AO)
- AO error budget
- Post-AO point-spread function morphology
- Anisoplanatic error study (ideal AO system)

# (IDL preliminaries...)

- launch IDL (or IDLDE=IDL+interface), with an OCA VPN running if doing it through the wifi.
- test it:  
IDL> 'hello'  
IDL> plot, [1,2], [1,2]




=> problems with the VPN ?

=> problems with the IDL graphical windows (plot) ?

# Images & turbulence - 1

The image formed through turbulent atmosphere (optically speaking) is degraded:

- Scintillation (due to intensity fluctuation in the pupil).
- Agitation (due to angle-of-arrival variation).
- Spreading (due to a loss of spatial coherence).



# Images & turbulence - 2


The object-image relation between the intensity  $I(\alpha)$  in the image plane (i.e. the focal plane of the telescope) and the brightness  $O(\alpha)$  of the object (in the sky) is a relation of convolution implying the point-spread function (PSF)  $S(\alpha)$  of the whole ensemble telescope+atmosphere, with  $\alpha$  the angular coordinates in the focal plane:

$$I(\vec{\alpha}) = O(\vec{\alpha}) * S(\vec{\alpha})$$


# Images & turbulence - 3

$$I(\vec{\alpha}) = O(\vec{\alpha}) * S(\vec{\alpha})$$


This relation is valid when the system is invariant by translation (i.e. everything happens within the isoplanatic domain)...



$\lambda/D$



$\lambda/r_0$






# Images & turbulence - 4




Some orders of magnitude concerning the turbulent atmosphere:

	$\lambda = 500 \text{ nm}$	$\lambda = 2.2 \mu\text{m}$
Fried parameter ( $r_0$ )	$\rightarrow 10 \text{ cm}$	60 cm
velocity of the turbulent layers ( $v$ )	$\rightarrow 10 \text{ m/s}$	id.
=> image FWHM ( $\epsilon \approx \lambda/r_0$ )	$\rightarrow 1''$	$\sim 1''$
=> evolution time ( $\tau_0 \propto r_0/v$ )	$\rightarrow 3 \text{ ms}$	18 ms

# Images & turbulence - 5



# Images & turbulence - 6



# Images & turbulence - 7

entrance pupil



image on the focal plane



remembering eq. 2.17 from  
the course of Éric Aristidi:

$$I(x, y) = \frac{1}{\lambda^2 F^2} \left| \hat{f}_0 \left( \frac{x}{\lambda F}, \frac{y}{\lambda F} \right) \right|^2$$

directly coming from (eq. 2.16):

$$f_F(x, y) = \frac{e^{ikF}}{i\lambda F} e^{\frac{i\pi\rho^2}{\lambda F}} \hat{f}_0 \left( \frac{x}{\lambda F}, \frac{y}{\lambda F} \right)$$

# Images & turbulence - 8

The wavefront is, modulo  $\lambda/2\pi$ , proportional to the phase  $\Phi(r)$  of the wave  $\Psi(r)$  which has went through the turbulent atmosphere before reaching the telescope:


$$\Psi(\vec{r}) = A(\vec{r}) \exp\{i\Phi(\vec{r})\}$$

Note that this phase can be decomposed following a base of polynomials, for example Zernike ones:

$$\Phi(\vec{r}) = \sum_i a_i Z_i(\vec{r})$$

# Images & turbulence - 9

## polynômes de Zernike



entrance pupil




image on the focal plane

# Images & turbulence - 10

turbulence intensity [ $m^{1/3}$ ]

$$r_0 = 0.185 \lambda^{5/6} \cos(\gamma)^{3/5} \left[ \int_0^\infty C_n^2(z) dz \right]^{-3/5}$$

dimension of  $r_0$  ? value in band H knowing  $r_0$  at 500nm (10cm) ?...

$$\tau = 0.36 \frac{r_0}{v}$$

$$\epsilon = 0.98 \frac{\lambda}{r_0}$$

$$\theta_0 = 0.314 \frac{r_0}{\bar{h}}$$

$$\bar{v} = \left( \frac{\int C_n^2(h) v(h)^{5/3} dh}{\int C_n^2(h) dh} \right)^{3/5}$$

$$N_s \simeq 0.34 \left( \frac{D}{r_0} \right)^2$$

$$\bar{h} = \left( \frac{\int C_n^2(h) h^{5/3} dh}{\int C_n^2(h) dh} \right)^{3/5}$$

Number of speckles for  $r_0=10\text{cm}$  and  $D=1\text{m}$  ?  
(also in H band)

# Images & turbulence - 11

$r_0$  in band H knowing  $r_0$  at 500nm ?...

$$r_0 = 0.185 \lambda^{\frac{6}{5}} \cos(\gamma)^{\frac{3}{5}} \left[ \int_0^\infty C_n^2(z) dz \right]^{-\frac{5}{3}}$$

$$r_0^{\text{H}=1.65 \mu\text{m}} = r_0^{500 \text{ nm}} \left( \frac{1.65}{0.5} \right)^{\frac{6}{5}} \simeq 0.42$$

Number of speckles for  $r_0=10\text{cm}$  and  $D=1\text{m}$  ?...

$$N_S^{500 \text{ nm}} \simeq 0.34 \left( \frac{1.0}{0.1} \right)^2 \simeq 34$$

$$N_S^{\text{H}} \simeq 0.34 \left( \frac{1.0}{0.42} \right)^2 \simeq 2$$

# Images & turbulence - 12

$$\Phi_{\varphi}(\vec{\nu}) = 0.0228 r_0^{-\frac{5}{3}} \left( \nu^2 + \frac{1}{\mathcal{L}_0^2} \right)^{-\frac{11}{6}}$$


Power Spectral Density (PSD) of the phase, function of the spatial frequency

## Kolmogorov/von Kármán model

- Kolmogorov : outerscale of turbulence  $\mathcal{L}_0$  is infinite.
- One can refine the model by considering also  $\ell_0$ .
- $\exists$  other models with a finite  $\mathcal{L}_0$  (and a non-zero  $\ell_0$ ).

Energy cascade: wind shear  $\Rightarrow$  turbulent energy injected into the system via a large eddy ( $\mathcal{L}_0$ ) which splits into smaller and smaller eddies ( $\ell_0$ ), and is finally viscously dissipated. Interval  $[\ell_0, \mathcal{L}_0] =$  inertial range.

# (A reminder of discrete Fourier transform (DFT)...)



# Images & turbulence - 13

$$\Phi_{\varphi}(\vec{\nu}) = 0.0228 r_0^{-\frac{5}{3}} \left( \nu^2 + \frac{1}{L_0^2} \right)^{-\frac{11}{6}}$$

Which, numerically written, and by considering wavefronts made of 'dim' pixels corresponding to 'L' meters, becomes:  
(re-writing - "de-dimensionalizing" - the equation with  $L_0=L_0 L/L$  and  $\nu=\nu L/L\dots$ )

```
freq = findgen(dim)
dsp   = .0228*(L/r0)^(5/3.)*L^2*(freq^2+(L/L0)^2)^(-11./6)
```

And which (with the right frequency scale) can be plot with:

```
plot_oo, 1./L*findgen(dim), dsp, XR=[1/L/1.2,dim*1/L*1.2], /XS
```

=> make a function that computes  $\text{PSD}(L_0, r_0, \text{dim}, L)$  and plot it for different  $[r_0, L_0] \dots$  [with, e.g:  $\text{dim}=1000L$ ,  $L=100.$ ,  $r_0=0.05..0.1..0.2$ ,  $L_0=100..10..1$ ]

# Images & turbulence - 14

Example of a function that computes the sum of two parameters:

```
function sum2par, par1, par2
  result=par1+par2
return, result
end
```

Compile and run the function (written, e.g., in a file sum2par.pro):

```
idl > .r sum2par
→ % Compiled module: SUM2PAR
idl > res = sum2par(2,1)
idl > res                                     (older version: idl> print, res)
→ 3
```

# (IDL preliminaries again...)

- test it more:

```
IDL> xx=findgen(50)
```

```
IDL> yy=2*xx+10
```

```
IDL> plot, xx, yy, TIT='my plot', XTIT='xx', YTIT='yy', $  
      XR=[-2,51], /XS, YR=[0,118], /YS
```

```
IDL> oplot, xx, min(yy)+max(yy)-yy, PS=4
```

